

The first X-ray eclipse of IGR J16479-4514?

E. Bozzo*, L. Stella*, G. Israel*, M. Falanga[†] and S. Campana**

*INAF - Osservatorio Astronomico di Roma, Via Frascati 33, 00044 Rome, Italy.

[†]CEA Saclay, DSM/IRFU/Service d'Astrophysique, F-91191, Gif sur Yvette, France.

**INAF - Osservatorio Astronomico di Brera, via Emilio Bianchi 46, I-23807 Merate (LC), Italy.

Abstract. We report on the first long (~ 32 ks) pointed *XMM-Newton* observation of the supergiant fast X-ray transient IGR J16479-4514. Results from the timing, spectral and spatial analysis of this observation show that the X-ray source IGR J16479-4514 underwent an episode of sudden obscuration, possibly an X-ray eclipse by the supergiant companion. We also found evidence for a soft X-ray extended halo around the source that is most readily interpreted as due to scattering by dust along the line of sight to IGR J16479-4514.

Keywords: X-rays: binaries - binaries: eclipsing - stars: individual (IGR J16479-4514)

PACS: 97.80.Jp

1. INTRODUCTION

Supergiant Fast X-ray transients (SFXTs) are a recently discovered subclass of high mass X-ray binaries. These sources display sporadic outbursts lasting from minutes to hours with peak luminosities of $\sim 10^{36}$ - 10^{37} erg s⁻¹, and spend long time intervals at lower X-ray luminosities, ranging from $\sim 10^{34}$ erg s⁻¹ to $\sim 10^{32}$ erg s⁻¹. Proposed models to interpret the SFXT behavior generally involve accretion onto a neutron star (NS) immersed in the clumpy wind of its supergiant companion [4]. The X-ray luminosity during lower activity states is likely due to accretion onto the NS at much reduced rate than in outburst [10, 9, 2]. Here we report on the first long, high sensitivity observation (~ 32 ks) of the SFXT IGR J16479-4514. This observation was carried out shortly after *Swift* discovered a very bright outburst from this source [8], and was aimed at investigating the low level emission of this source and gaining insight in the physical mechanisms that drive this activity.

2. DATA ANALYSIS AND RESULTS

XMM-Newton [5] observed IGR J16479-4514 between March 21 14:40:00UT and March 22 01:30:00UT. We show in figure 1 the 2-10 keV EPIC-PN light curve and spectra of the observation. In the first ~ 4 ks IGR J16479-4514 was caught during the decay from a higher (“A”) to a lower (“B”) flux state. The shape of the above light curves and evolution of the spectrum across the state A-state B transition presented remarkable similarities with the eclipse ingress of eclipsing X-ray sources such as OAO 1657-415 [1]. Moreover the slower decay of the soft X-ray light curve (in turn similar to that observed in OAO 1657-415) suggested that IGR J16479-4514 is seen through an extended dust-scattering halo [3]. To confirm this we analysed the radial distribution of the X-ray




FIGURE 1. *Left panel:* EPIC-PN light curve of IGR J16479-4514 in the 2-10 keV band. The bin time is 100 s and the start time is March 21 16:42:42UT. *Right panel:* 2-10 keV spectra extracted from intervals A and B. The best fit models and the residuals from these fits are also shown. The best fit parameters for state A ($\chi^2/\text{d.o.f.} = 20.4/26$) are: $N_{H,\alpha} = 35^{18}_{-13} \times 10^{22} \text{ cm}^{-2}$, $N_{H,\beta} = 9^{9}_{-6} \times 10^{22} \text{ cm}^{-2}$, $I_{\alpha} = 1.8^{0.4}_{-0.3} \times 10^{-3}$, $I_{\beta} = 5^{8}_{-4} \times 10^{-3}$, $E_{\text{ln}1} = 6.53^{0.06}_{-0.07} \text{ keV}$, $I_{\text{ln}1} = 5^{3}_{-3} \times 10^{-5}$, $EW_{\text{ln}1} = 0.15 \text{ keV}$. The measured X-ray flux in this state is $F_{2-10 \text{ keV}} = 10^{-11} \text{ erg/cm}^2/\text{s}$, corresponding to $L_X = 2.9 \times 10^{34} \text{ erg/s}$ [the source distance is 4.9 kpc; 6]. For the B state ($\chi^2/\text{d.o.f.} = 36.5/33$) we found: $N_{H,\alpha} = 54^{26}_{-25} \times 10^{22} \text{ cm}^{-2}$, $I_{\alpha} = 1.1^{0.5}_{-0.4} \times 10^{-4}$, $I_{\beta} = 8^{8}_{-3} \times 10^{-4}$, $E_{\text{ln}1} = 6.51^{0.03}_{-0.02} \text{ keV}$, $I_{\text{ln}1} = 1.8^{1.0}_{-0.4} \times 10^{-5}$, $EW_{\text{ln}1} = 0.77 \text{ keV}$, $E_{\text{ln}2} = 7.11^{0.06}_{-0.09} \text{ keV}$, $I_{\text{ln}2} = 8^{7}_{-4} \times 10^{-6}$, $EW_{\text{ln}2} = 0.28 \text{ keV}$, $F_{2-10 \text{ keV}} = 7.5 \times 10^{-13} \text{ erg/cm}^2/\text{s}$ ($L_X = 2.2 \times 10^{33} \text{ erg/s}$). Here $F_{2-10 \text{ keV}}$ is the absorbed flux in the 2-10 keV band, and errors are at 90% confidence level.

photons detected from IGR J16479-4514 and compared it with the point spread function (PSF) of the *XMM-Newton* telescope/EPIC-PN camera. We found that a 30'' extended dust scattering halo located halfway between us and IGR J16479-4514 agrees well with the e-folding decay time of the soft X-ray light curves. Motivated by the above findings we considered a spectral model that has been used in studies of eclipsing X-ray binaries seen through a dust-scattering halo, that is: $I(E) = e^{\sigma(E)N_{H,\beta}} I_{\beta} E^{-\beta} + e^{\sigma(E)N_{H,\alpha}} [I_{\alpha} E^{-\alpha} + I_{\text{ln}1} e^{-(E-E_{\text{ln}1})^2/(2\sigma_{\text{ln}1}^2)} + I_{\text{ln}2} e^{-(E-E_{\text{ln}2})^2/(2\sigma_{\text{ln}2}^2)}]$. In practice we fitted a model with two continuum components, a single power law of slope α (with normalisation I_{α} and column density $N_{H,\alpha}$) and a power law of slope β (with normalisation I_{β} and column density $N_{H,\beta}$). The former component was taken to approximate the sum of two α -slope power laws: the first represents the direct component dominating before the eclipse ingress (i.e. the component due to the accretion process onto the NS), whereas the second represents the wind scattered component dominating during the eclipse [1]. The second power law component, with photon index $\beta = \alpha + 2$, originates from small angle scattering of (mainly) direct photons off interstellar dust grains along the line of sight [3]. The Gaussians in the above equation represent iron features around $\sim 6.4 \text{ keV}$ and $\sim 7.0 \text{ keV}$, arising from the reprocessing of the source radiation in the supergiant wind. We kept α fixed at 0.98, and thus $\beta = 2.98$ [7]. The best fit parameters are reported in the caption of Fig. 1.

In state A the relatively high source flux and small EW of the iron fluorescence line testifies that (most of) the emission is likely due to the direct component. In state B the ratio I_{α}/I_{β} is larger than the corresponding value obtained during state A, whereas the EW of the Fe-line at $\sim 6.5 \text{ keV}$ increased from $\sim 150 \text{ eV}$ to $\sim 770 \text{ eV}$. We also found




FIGURE 2. Simulated Simbol-X spectrum of IGR J16479-4514 during state B. Both the results for the MPD (Macro-Pixel Detector), and the CZT(Cd(Zn)Te detector) are shown.

evidence ($\sim 2\sigma$) for an additional Fe-line at ~ 7.1 keV with a ~ 300 eV EW, consistent with being the K_β . The most natural interpretation of this is that in state B the direct emission component is occulted along our line of sight, while the spectrum we observe is the sum of a dust scattered component, dominating at lower energies, and a wind scattered component characterised by a high absorption. The marked increase in the EW of the Fe-line at ~ 6.5 keV across the state A-state B transition testifies that the region where the line is emitted is larger than the occulting body (the supergiant companion, if we are dealing with an eclipse); this also provides further evidence that the unocculted emission in state B at hard X-ray energies (the α -slope power law) arises mostly from photons scattered by the wind in the immediate surrounding of the source.

We conclude that IGR J16479-4514 is the first SFXT that displayed evidence for an X-ray eclipse. Further observations of IGR J16479-4514 and other SFXTs in quiescence, will improve our knowledge of the low level emission of these sources, and will help clarifying if X-ray eclipses are common in SFXTs. In Fig. 2 we show the X-ray spectrum of IGR J16479-4514 during state B that would be observed by Simbol-X. We used an exposure time of 28 ks and the latest Simbol-X simulation tool available (see <http://www.apc.univ-paris7.fr/Simbolx2008/>).

We thank Norbert Schartel and the XMM-Newton staff, for carrying out this ToO observation; EB thanks L. Sidoli and P. Romano for their collaboration during the early stages of this work.

REFERENCES

1. Audley, M.D., Nagase, F., Mitsuda, K., Angelini, L., Kelley, R.L. 2006, MNRAS, 367, 1147
2. Bozzo, E., Falanga M., Stella L. 2008, ApJ, 683, 1031
3. Day, C.S.R. & Tennant, A.F. 1991, MNRAS, 251, 76
4. in 't Zand 2005, A&A, 441, L1
5. Jansen, A., Lumb, D., Altieri B., et al. 2001, A&A, 365, L1
6. Rahoui, F., Chaty, S., Lagage, P., Pantin, E. 2008, preprint (astro-ph/0802.1770)
7. Romano, P., et al. 2008a, ApJ, 680, L137
8. Romano, P., et al. 2008b, Astr. Tel., 1435
9. Sidoli, L., et al. 2007, A&A, 476, 1307
10. Walter, R. & Zurita Heras, J. A. 2007, A&A, 476, 335